#### **HOW TO TAKE**

# **PLENVU®**



## AFTERNOON APPOINTMENT

For full details, please refer to the patient information leaflet (PIL) supplied with your box of Plenvu.

#### Reporting of side effects in the United Kingdom

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at https://yellowcard.mhra.gov.uk/.

By reporting side effects you can help provide more information on the safety of this medicine.

Norgine Adverse events should also be reported to

**Norgine** Adverse events should also be reported to Medical Information at Norgine Pharmaceuticals on +44 1895 826606 or E-mail: medinfo@norgine.com

#### Reporting of side effects in Ireland

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly to the HPRA, via www.hpra.le, medsafety@hpra.le or by calling (01) 676 4971. By reporting side effects you can help provide more information on the safety of this medicine. Norgine Adverse events should also be reported to Medical Information at Norgine Pharmaceuticals on +44 1895 826606 or E-mail: medinfo@norgine.com

For more information please visit

### bowelcleansing matters.co.uk





#### **CLEAR FLUIDS**

Remember you can keep drinking until 2 hours before your colonoscopy.



Water



Diluted cordials (NOT blackcurrant)



Herbal tea



Clear soup



Clear fruit juice (without pulp)



Black tea/coffee (without milk)

#### EATING SUGGESTIONS

Your healthcare professional may request that you adopt a low fibre diet when preparing for your bowel preparation.





For specialised diets, please refer to your healthcare professional.



## Unless your doctor or nurse tells you differently, start taking Plenvu as outlined below.

For full details, please refer to the patient information leaflet (PIL) supplied with your box of Plenvu.

## All sachets must be taken with the right amount of water according to the timings below.

You will have lots of watery bowel movement, so **stay near a toilet** because you may need to use it urgently.

#### DAY OF colonoscopy DAY BEFORE colonoscopy Afternoon Night Morning Afternoon Evening Morning Stop all intake Take Take Stop all intake of DOSE 2 of food\* DOSE 1 fluids 2 hours 3pm 6pm 6am before colonoscopy Drink Drink If thirsty, Extra clear Light Light 500ml of drink plenty 500ml of fluids\* ••• breakfast\* lunch\* clear of clear clear only fluids\* fluids\* fluids\*

### DOSE 1 6pm



Open the carton and remove the **Dose 1** sachet.



Pour the contents of the **Dose 1** sachet into a measuring container that can hold at least 500ml of fluid.



Add water to make up to 500ml and stir until all the powder has fully dissolved.



Each dose of Plenvu must be **sipped slowly** over 30 minutes, followed by 500ml of mandatory clear fluid over a further 30 minutes.

### DOSE 2 6am



Open the carton and remove the **Dose 2** sachets A and B.

Overnight



Add water to make up to 500ml and stir until all the powder has fully dissolved.



Pour the contents of **Dose 2** sachets A and B into a measuring container that can hold at least 500ml of fluid.



Each dose of Plenvu must be **sipped slowly** over 30 minutes, followed by 500ml of mandatory clear fluid over a further 30 minutes.

#### **Hints & Tips**



**Sip slowly** over at least 30 minutes



Once dissolved, (this may take up to 8 minutes) the solution can be refrigerated



Try using a straw



Stay hydrated with plenty of clear fluids\*

<sup>\*</sup> Please see overleaf.